

## OUTCOME OF PEDIATRIC MISSILE HEAD INJURIES

HUSSEIN IMRAN MOUSA

Neurosurgical Department Alsadr Teaching Hospital, Basrah, Iraq

### ABSTRACT

#### Introduction

Missile injuries have progressively increased in Iraq after 2003. These injuries are uncommon in children, but cause greater severity compared to other injuries.

#### Method

Fifty two patients with pediatric missile injuries (PMIs) admitted to Alsadre teaching hospital (Basrah neurosurgical center) from March 2006 to June 2014 were enrolled in this study.

#### Results

The patients' age ranged from 6 months-18 years with a male/ female ratio= 11:2. The patients were victims of roadside grenade attacks, terror explosions, haphazard fires between security forces and militants, or air firing on marriages and processions ceremony. Bullet injury accounted for 44% of the study group, whereas the rest of the PMIs were caused by fragments from blast injuries. In patients with a Glasgow Coma Score (GCS) of 3–8 there was a 50% mortality rate and none of the patients had a good outcome. In patients with a GCS score of 9–12 there was no mortality rate and 60% of the patients had a good outcome. There was one death in the group with GCS scores of 13-15. Pupillary abnormalities were seen in 7 cases. About 73% of the patients had a unihemispheric injury, 13.5% had intracerebral hemorrhage, 9.6% had midline shift in addition to a unihemispheric injury, whereas 3.8% had bihemispheric involvement. About 15.4% of the patients had motor weakness. Two of the motor weakness and 3 of non-motor weakness patients died. Sixteen patients (30.8%) had fits. Four patients of them died, while one patient died from the group who did not develop fits. Four patients (7.7%) had meningitis. Thirty-four patients (65.4%) had some psychological upsets.

#### Conclusions

PMIs do not differ significantly from other types of head injury as far as prognostic factors are concerned. In Iraq, the epidemiology of PMI is different from that in the West. Education, of the community on the risks of haphazard bullet injuries, is recommended and the government must show more power to avoid these wrong habits.

**KEYWORDS:** Pediatric Missile Injuries, Bad Outcome and Good Outcome

### INTRODUCTION

- Missile-related injuries form a leading cause of death and these injuries have progressively increased in Iraq after 2003.<sup>1-2</sup>
- Missile injuries are uncommon in children, but cause greater severity, need for major surgery, mortality, disability and costs compared to other types of injury. Although, they are uncommon, there is a large variation in the incidence of pediatric missile injuries between regions.<sup>3</sup>

- Pediatric missile injury has been the subject of several population-based epidemiologic studies from the mid-1980s through 2002.<sup>4-8</sup> More recent population-based research has been limited and focused exclusively on mortality<sup>9-12</sup> or was restricted to hospital-based samples.<sup>13-15</sup> Data from the National Violent Death Reporting System provide insight into the problem.<sup>16</sup>

## METHODS

Eighteen year-old or younger fifty-two patients, with PMIs due to a bullet or fragments from a blast whom were admitted to Alsadre Teaching Hospital (the only Basrah Neurosurgical Center) between March 2006 and June 2014, were enrolled in this study regardless their neurological status.

Data were prospectively collected from 2006 onwards. Patients with associated injuries, such as those of the chest, abdomen, spine, and long bones, which could contribute to a low neurological status by causing hemorrhagic shock or hypoxia, were excluded. This exclusion was considered because we wanted to study only the aspect of head injury caused by bullets or blast fragments.

Most of the patients were transported by civilians; the interval between injury and admission varied from 30 to 180 minutes depending on the distance of the site of injury from the Centre and direct or shifting from other hospitals.

Resuscitation was performed to all of the patients according to the advanced trauma life support guidelines; the nature of the resuscitation depended upon the clinical status of the patient. After resuscitation, patients were assessed neurologically and those with a GCS score  $\leq 8$  were intubated (a postresuscitation GCS score was used in further analysis). A head CT scan was obtained in all cases. After CT, patients were surgically treated depending on the nature of the injury(s).

Surgery consisted of decompressive craniectomy and removal of necrotic brain, hematoma, debris, and bone fragments that were easily accessible. No attempt was made to perform extensive debridement of the missile tract in the brain. Firstly, superficial debridement and then duraplasty and scalp closure were performed.

All the 52 patients received antibiotics (combination of ceftriaxone + ampiclox) along with phenytoin 5 mg/kg/day after administering a loading dose of 15 mg/kg. Osmotherapy with mannitol (1 g/kg/day in divided doses) was delivered to all the patients who had brain edema. Intracranial pressure monitoring was not done because of the unavailability of the equipment.

All surgeries were performed by neurosurgeons and not by trauma surgeons. Outcome was followed up using the Glasgow Coma Scale from 6 months-6 years. The bad outcome groups included grades were as follows: 5= death, 4= vegetative state, 3= severe disability but conscious and a good outcome groups: 2= minimal disability, and 1= good recovery.

All data were analyzed using SPSS 19 software. Predictor of outcome were assessed using chi-square analysis, and then logistic regression analysis was used, a p-value  $< 0.05$  taken as significant.

## RESULTS

The demographic characteristics: The 52 patients included in our study ranged in age from 2 to 18 years, with a mean age of 10.1 years. There was no significant statistical association could be recognized between age and outcome ( $p = 0.13$ ). The male/female ratio was 11:2 with a high male preponderance.

Mechanism of injury: all the children were victims of violence in which they had no role and victims of roadside terror explosions, haphazard fire of between security forces and militants or firing in the air as a habit after winning a football match, marriages and processions ceremony. Some children were injured while playing with unidentified objects found in grazing fields, which turned out to be explosives. There were no instances of suicide or murder among the study population. Bullet injuries accounted for 44% of the study group, whereas the rest of the PMIs were caused by fragments from blast injuries. There was no significant difference in outcome between the two groups ( $p = 0.15$ ).

### Clinical Factors

Glasgow Coma Score: In patients with a score of 3–8 there was a 50% mortality rate and none of the patients had a good outcome. In patients with a GCS score of 9–12 there was no mortality and 60% of the patients had a good outcome. There was one death in the group with GCS scores of 13–15 (sudden onset of severe headache followed by status epilepticus and Brain CT scan showed multiple intraparenchymal hemorrhage) otherwise the other patients had a good outcome ( $p < 0.001$ ).

### Pupillary Status

Pupillary abnormalities were seen in 7 cases: anisocoria in 3 cases (5.8%) and dilated nonreactive pupils in 4 cases (7.7%). All patients in the latter group died, while there was no mortality in the anisocoria group, but all patients had a bad outcome. One patient died with normal pupils but the other 84.3% survived ( $p < 0.001$ ).

Radiological Parameters: Computed tomography scanning done to all the study population; 38 patients (73%) had a uni-hemispheric injury, 7 patients (13.5%) had intracerebral hemorrhage and 5 patients (9.6%) had midline shift in addition to a uni-hemispheric injury, whereas 2 (3.8%) had bi-hemispheric involvement. The bi-hemispheric involvement group followed by the intracerebral hemorrhage with midline shift group had significantly bad outcome as compared to the uni-hemispheric injury group ( $p = 0.017$ ).

**Motor Weakness:** There were 8 patients (15.4%) had motor weakness while 44 patients (84.6%) had no weakness. Two (25%) of the motor weakness patients died while 3 (6.8%) of non-motor weakness patients died ( $p = 0.001$ ).

### Complications

**Seizure:** Seizures had in 19% of patients with bad outcome groups as compared to 5.8% of patients in a good outcome groups ( $p = 0.001$ ).

**Meningitis:** Four (7.7%) patients had meningitis. All of them treated and became well ( $p = 0.45$ ).

**Hydrocephaly:** Three patients (5.8%) had hydrocephaly. All of the patients underwent ventriculo-peritoneal shunt. In one of them, external ventricular drain was done firstly, then after 7 months hydrocephaly developed and ventriculo-peritoneal shunt was done. There were no significant outcomes in patients who had hydrocephaly as compared to those none had hydrocephaly ( $p = 0.08$ ).

**Psychological Upset:** Thirty four patients (65.4%) had some psychological upset. All bad outcome groups (13 patients) had psychological upset while 21(40%) patients from a good outcome groups had psychological upset ( $p=0.02$ ).

## DISCUSSIONS

In studies from more developed countries, the most common cause of PMIs in children is homicide or suicide. Among the Iraqi population, there are some incidents occur to people being targeted by explosive device blasts or of bombs being planted in public places by terrorists. People in the present study were also injured during firing in the air as a habit after winning a football match, marriages and processions ceremony. Since the grenade attacks were carried out on security forces in the vicinity of residential quarters and market areas, there were cases of injuries caused by grenade fragments, which otherwise would not have been expected to occur in a civilian population. None of our patients reported lying down on hearing missiles or after a blast; all of them had run for safe shelter, which made them more susceptible to stray bullets. Perhaps this tendency to run explains the higher incidence of head injury in these children. If people are taught to lie down in these circumstances, then there may be a decrease in the incidence of head injury. Note that patients who were injured by bomb blasts and improvised explosive device blasts were excluded from this study because they had associated injuries other than head trauma.<sup>39,40,41,42,43,44</sup>

Age has been shown to have a varying effect on the outcome in different series. In some studies, with decreasing age, outcome was adversely affected,<sup>35,19,20,36,37,38</sup> while in others, it was the opposite.<sup>11</sup> In this study no age group had a better outcome ( $p=0.13$ ). This reflected haphazard of PMI in the population.

By comparing mechanisms of injury, most of the reference bullet injury carry bad outcome than blast injury.<sup>47,48,32,33</sup> In this study there were no differences in the outcome in patients with bullet injuries as compared to the patients with injuries from grenade blast fragments ( $p=0.15$ ).; a good outcome occurred in 34.6% of patients in the first group as compared to 40.4% in the latter. The difference documented in other studies may be the result of the fact that PMIs in children is homicide or suicide while accidental injury with far distances causes the bullet losses its high energy ( $E = \frac{1}{2}mv^2$ ). This is same causes lead to the mortality rate 17.64% lower the others studies. Nelson and colleagues,<sup>56</sup> the mortality rate was 36%. In Kaufman's study,<sup>22</sup> the survival rate was only 2%.

Sixteen patients (30%) with severely GCS at time of admission to hospital all of them had bad outcome as compared with 4 patients with moderate GCS had bad outcome and 1 patient with mild GCS died. The GCS score was documented to be the strongest predictor of outcome after head injury,<sup>17-27</sup> and this trend was seen in the current study group as well  $p$  value 0.001.

None of this study patient with fixed dilated pupils survived, and no patients with anisocoria had a good outcome as compared to 71% of the patients with normal pupils. Patients with normally reacting pupils had a good outcome than those with nonreactive pupils. Pupillary changes reflect the extent of brainstem compression, which has a significant effect on outcome.<sup>29,31</sup> A unilateral fixed dilated pupil is thought to be attributable to the third nerve compression, while bilateral dilation signifies disturbed brainstem function.<sup>17,21,32,33,34</sup>

In the bihemispheric injury, midline shift, and intracerebral hemorrhage had a good outcome 1.9%, 3.8%, and 5.8% respectively as compared to 63.5% in the unilateral injury group. This trend is obviously due to involvement of the critical midline structures. Eisenberg et al.<sup>45</sup> reported that midline shift is a very strong predictor of abnormal intracranial

pressure, and the risk of death is greater if the midline shift is large. Conversely, the investigations of Miller et al. demonstrated no or only a poor correlation between midline shift and intracranial pressure<sup>46</sup>. In intracerebral hemorrhage more damage to the brain results as primary injury and/or swelling of brain due to edema causes damage to the neuron as a secondary injury. In this study there was a significant correlation between the bihemispheric injury, midline shift, and intracerebral hemorrhage and bad outcome. P value 0.017

### Complications of PMI

Seizures occurred in 19% of patients with bad outcome groups as compared to 5.8% of patients in a good outcome groups, so the fit is a statically significant factor ( $p=0.001$ ). Seizures indicated injury to brain parenchyma and the formation of gliotic scar tissue in the tract of the missile injury and considered as a risk factor in the most of the study sample.

Meningitis occurred in 4 patients; three patients in a good outcome groups and one patient in the bad outcome groups. In this study meningitis has not an additional risk factor for prognosis ( $p=0.46$ ). This might be due to the low number of cases and its treatable disease. This is same causes lead to a hydrocephaly not a significant factor for prognosis ( $p=0.08$ ) when it occurred in 3 patients, two patients in a good outcome groups and one patient in the bad outcome groups. Kishor et al reported its incidence to be up to 15% among all patients with severe head injury. According to Zander and Foroglou<sup>2</sup>, about 10% of patients in large sample of head injured patients with prolonged coma developed some degree of hydrocephalus.<sup>55</sup>

The wide spectrum of post-traumatic sequelae presented by the victim of craniocerebral injury is impressive, and ranges from a comatose, purely vegetative akinetic mutism to neurological syndromes of varying severity and mild personality disorders. The main organic cause of disability after a head injury is disturbance of mental capacity. This disturbance may occur at various levels of mental activity and may range from affection of well-established skills such as speech, reading, calculation and orientation, to disturbances of higher capacities such as memory, abstract

Thinking and reasoning.<sup>49-55</sup> In this study 34 patients (65.4%) had psychological upset included all bad outcome groups (13 patients) and 21 (40%) patients from a good outcome groups. There were psychological upset considered statically a significant factor for bad prognosis ( $p=0.02$ ). This might be due to wide damaging neurons and brain concussion when fall on ground after injuries.

### CONCLUSIONS

- PMIs do not differ significantly from head injury in other age groups as far as prognostic factors are concerned.
- The GCS score was documented to be the strongest predictor of outcome after head injury and can add three other strong predictors of outcome in could be diagnosed, these were pupillary abnormality, motor deficit and seizures.
- In the developing countries, the epidemiology of PMI is different from that in the West. Education of the community to word risk of haphazard bullet injuries is needed and the government must be more powerful to avoid these wrong habits.

### REFERENCES

1. Zouris JM, Walker GJ, Dye J, Galarneau M. Wounding patterns for US marines and sailors during Operation Iraqi

- Freedom, major combat phase. *Mil Med.* 2006;171:246–52. [PubMed]
2. Chambers LW, Rhee P, Baker BC, Perciballi J, Cubano M, Compeggie M, et al: Initial experience of US Marine Corps forward resuscitative surgical system during Operation Iraqi Freedom. *Arch Surg* 140:26–32, 2005
  3. DiScala C, Sege R. Outcomes in children and young adults who are hospitalized for firearms-related injuries. *Pediatrics.* 2004; 113(5):1306–1312
  4. Nance ML, Denysenko L, Durbin DR, BranasCC, Stafford PW, Schwab CW. The ruralurbancontinuum: variability in statewideserious firearm injuries in children andadolescents. *Arch PediatrAdolesc Med.* 2002;156(8):781–785
  5. Powell EC, Jovtis E, Tanz RR. Incidence andcircumstances of nonfatal firearm-relatedinjuries among children and adolescents. *Arch PediatrAdolesc Med.* 2001;155(12):1364–1368
  6. Roesler J, Ostercamp M. Pediatric firearm injury in Minnesota, 1998. Fatal and nonfatalfirearm injuries among Minnesotayouth.*Minn Med.* 2000;83(9):57–60
  7. Zavoski RW, Lapidus GD, Lerer TJ, Banco LI.A population-based study of severe firearm injury among children and youth.*Pediatrics.*1995;96(2 pt 1):278–282
  8. Dowd MD, Knapp JF, Fitzmaurice LS. Pediatricfirearm injuries, Kansas City, 1992:a population-based study. *Pediatrics.* 1994;94(6 pt 1):867–873
  9. Sege RD, Kharasch S, Perron C, et al. Pediatricviolence-related injuries in Boston:results of a city-wide emergency departmentsurveillance program. *Arch PediatrAdolesc Med.* 2002;156(1):73–76
  10. Cheng TL, Wright JL, Fields CB, et al. Violentinjuries among adolescents: declining morbidityand mortality in an urban population. *Ann Emerg Med.* 2001;37(3):292–300
  11. Nance ML, Carr BG, Kallan MJ, Branas CC,Wiebe DJ. Variation in pediatric and adolescentfirearm mortality rates in rural andurban US counties. *Pediatrics.* 2010;125(6):1112–1118
  12. Kegler SR, Annest JL, Kresnow M, Mercy J; Centers for Disease Control and Prevention(CDC). Violence-related firearm deathsamong residents of metropolitan areas andcities—United States, 2006–2007. *MMWRMorb Mortal Wkly Rep.* 2011;60(18):573–578
  13. Martin CA, Unni P, Landman MP, et al. Racedisparities in firearm injuries and outcomesamong Tennessee children. *JPediatr Surg.* 2012;47(6):1196–1203
  14. Schecter SC, Betts J, Schecter WP, VictorinoGP. Pediatric penetrating trauma: the epidemiccontinues. *J Trauma Acute CareSurg.* 2012;73:721–725
  15. Senger C, Keijzer R, Smith G, MuenstererOJ. Pediatric firearm injuries: a 10-yearsingl-center experience of 194 patients. *JPediatr Surg.* 2011;46(5):927–932

19. Hernesniemi J: Penetrating craniocerebral missile wounds in civilians. *Acta Neurochir (Wien)* 49:199–205, 1979
20. Bruce DA, Raphaely RC, Goldberg AI, Zimmerman RA, Bilaniuk LT, Schut L, et al: Pathophysiology, treatment and outcome following severe head injury in children. *Childs Brain* 5:174–191, 1979
21. Bruce DA, Schut L, Bruno LA, Wood JH, Sutton LN: Outcome following severe head injuries in children. *J Neurosurg* 48:679–688, 1978
22. Humphreys RP: Outcome of severe head injury in children, in Raimondi AJ (ed): *Concepts in Pediatric Neurosurgery*. Basel: Karger, 1983, Vol 3, pp 191–201
23. Cernak I, Wang Z, Jiang J, Bian X, Savic J: Ultrastructural and functional characteristics of blast injury-induced neurotrauma. *J Trauma* 50:695–706, 2001
24. Krieger D, Adams HP, Schwarz S, Rieke K, Aschoff A, Hacke W: Prognostic and clinical relevance of pupillary responses, intracranial pressure monitoring, and brainstem auditory evoked potentials in comatose patients with acute supratentorial mass lesions. *Crit Care Med* 21:1944–1950, 1993
25. Langfitt TW, Gennarelli TA: Can the outcome from head injury be improved? *J Neurosurg* 56:19–25, 1982
26. Levy ML: Outcome prediction following penetrating craniocerebral
27. injury in a civilian population: aggressive surgical management in patients with admission Glasgow Coma Scale scores of 6 to 15. *Neurosurg Focus* 8(1):e2, 2000
28. Mayer T, Walker ML: Emergency intracranial pressure monitoring
29. in pediatrics: management of the acute coma of brain insult. *Clin Pediatr (Phila)* 21:391–396, 1982
30. Miller JD, Butterworth JF, Gudeman SK, Faulkner JE, Choi SC, Selhorst JB, et al: Further experience in the management of severe head injury. *J Neurosurg* 54:289–299, 1981
31. Sherman WD, Apuzzo ML, Heiden JS, Petersons VT, Weiss MH: Missile wounds to the brain—a civilian experience. *West J Med* 132:99–105, 1980
32. Taylor A, Butt W, Rosenfeld J, Shann F, Ditchfield M, Lewis E, et al: A randomized trial of very early decompressive craniectomy in children with traumatic brain injury and sustained intracranial hypertension. *Childs Nerv Syst* 17:154–162, 2001
33. Clusmann H, Schaller C, Schramm J: Fixed and dilated pupils after trauma, stroke, and previous intracranial surgery: management and outcome. *J Neurol Neurosurg Psychiatry* 71: 175–181, 2001
34. Cominos SC: Early prognosis of severe head injuries in children.
35. *Acta Neurochir Suppl (Wien)* 28:144–147, 1979
36. Zuccarello M, Facco E, Zampieri P, Zanardi L, Andrioli GC: Severe head injury in children: early prognosis and outcome. *Childs Nerv Syst* 1:158–162, 1985
37. Ropper AH: The opposite pupil in herniation. *Neurology* 40:1707–1709, 1990
32. Mazza C, Pasqualin A, Feriotti G, DaPian R: Traumatic extradural haematomas in children: experience with 62 cases. *Acta Neurochir (Wien)*

65:67–80, 1982

38. Liebenberg WA, Demetriades AK, Hankins M, Hardwidge C, Hartzenberg BH: Penetrating civilian craniocerebralmissile wounds: a protocol of delayed surgery. *Neurosurgery* 57:293–299, 2005
39. Michaud LJ, Rivara FP, Grady MS, Reay DT: Predictors of survival and severity of disability after severe brain injury in children. *Neurosurgery* 31:254–264, 1992
40. Luerssen T, Klauber MR, Marshall LF: Outcome from head injury related to patient's age. A longitudinal prospective study of adult and pediatric head injury. *J Neurosurg* 68:409–416, 1988
41. DePalma RG, Burris DG, Champion HR, Hodgson MJ: Blast injuries. *N Engl J Med* 352:1335–1342, 2005
42. Dhellemmes P, Lejeune JP, Christiaens JL, Combelles G: Traumatic
43. extradural hematomas in infancy and childhood. Experience with 144 cases. *J Neurosurg* 62:861–864, 1985
44. Pfenninger J, Kaiser G, Lütschg J, Sutter M: Treatment and outcome of the severely head injured child. *Intensive Care Med* 9:13–16, 1983
45. Raimondi AJ, Samuelson GH: Craniocerebralmissile wounds in civilian practice. *J Neurosurg* 32:647–653, 1970
46. Eisenberg HM, Gary HE Jr, Aldrich EF, Saydjari C, Turner B, Foulkes MA, et al: Initial CT findings in 753 patients with severe head injury. A report from the NIH Traumatic Coma Data Bank. *J Neurosurg* 73:688–698, 1990
47. Gutierrez FA, McLone DG, Raimondi AJ: Epidural hematomas in infancy and childhood, in American Society for Pediatric Neurosurgery (eds): *Concepts in Pediatric Neurosurgery*. Basel: Karger, 1981, Vol 1, pp 188–201
48. Jagannathan J, Okonkwo DO, Dumont AS, Ahmed H, Bahari A, Prevedello DM, et al: Outcome following decompressivetranscranial craniectomy in children with severe traumatic brain injury: a 10-year single-center experience with long-term follow up. *J Neurosurg* 106 (4 Suppl):268–275, 2007
49. Nelson TJ, Wall DB, Stedje-Larsen ET, Clark RT, Chambers LW, Bohman HR: Predictors of mortality in close proximity blast injuries during Operation Iraqi Freedom. *J Am Coll Surg* 202:418–422, 2006
50. Peleg K, Aharonson-Daniel L, Michael M, ShapiraSC : Patterns of injury in hospitalized terrorist victims. *Am J Emerg Med* 21:258–262, 2003
51. Rodoplu U, Arnold JL, Tokyay R, Ersoy G, Cetiner S, Yücel T: Mass-casualty terrorist bombings in Istanbul, Turkey, November 2003: 45. Raimondi AJ, Hirschauer J: Head injury in the infant and toddler.
52. Coma scoring and outcome scale. *Childs Brain* 11:12– 35, 1984
53. Miller JD, Becker DP, Ward JD, Sullivan HG, Adams WE, Rosner MJ: Significance of intracranial hypertension in severe head injury. *J Neurosurg* 47:503–516, 1977
47. Jandial R, Reichwage B, Levy M, Duenas V, Sturdivan L: Ballistics
54. for the neurosurgeon. *Neurosurgery* 62:472–480, 2008
55. Jennett B, Teasdale G, Braakman R, Minderhoud J, Knill- Jones R: Predicting outcome in individual patients after



- severe head injury. *Lancet* 1:1031–1034, 1976
56. Kluger Y: Bomb explosions in acts of terrorism—detonation, wound ballistics, triage and medical concerns. *Isr Med Assoc J* 5:235–240, 2003
  57. Kaufman HH: Civilian missile wounds to the head. *Neurosurgery* 32:962–964, 1993
  59. Kaufman HH, Levy ML, Stone JL, Masri LS, Lichtor T, Lavine SD, et al: Patients with Glasgow Coma Scale 3, 4, 5 after missile wounds to the brain. *NeurosurgClin N Am* 6: 701–714, 1995
  60. Hubschmann O, Shapiro K, Baden M, Shulman K: Craniocerebral missile injuries in civilian practice—prognostic criteria and surgical management: experience with 82 cases. *J Trauma* 19:6–12, 1979
  62. Lillard PL: Five year experience with penetrating craniocerebral missile wounds. *SurgNeurol* 9:79–83, 1978
  53. Miner ME, Cabrera JA, Ford E, Ewing-Cobbs L, Amling J: Intracranial penetration due to BB air rifle injuries. *Neurosurgery* 19:952–954, 1986
  64. 57. Kishor PRS, Lipper MH, Miller JD, Girevendulis AK, Becker DP, Vines FS: Post-traumatic hydrocephalus in patients with severe head injury. *Neuroradiology* 1978; 16:261-5.
  65. Kaufman HH, Schwab K, Salazar AM: A national survey of neurosurgical care for penetrating head injury. *SurgNeurol* 36:370–377, 1991
  66. Kunze E, Meixensberger J, Janka M, Sörensen N, Roosen K: Decompressive craniectomy in patients with uncontrollable intracranial hypertension. *Acta Neurochir Suppl* 71:16–18, 1998
  68. Levy ML, Masri LS, Lavine S, Apuzzo MLJ: Outcome prediction after penetrating craniocerebral injury in a civilian population: aggressive surgical management in patients with admission Glasgow Coma Scale scores of 3, 4, or 5. *Neurosurgery* 35:77–85, 1994
  70. Levy ML, Masri LS, Levy KM, Johnson FL, Martin-Thomson E, Couldwell WT, et al: Penetrating craniocerebral injury resultant from missile wounds: gang-related injury in children and adolescents. *Neurosurgery* 33:1018–1025, 1993
  71. Polin RS, Shaffrey ME, Bogaev CA, Tisdale NRN, Germanson T, Bocchicchio B, et al: Decompressive bifrontal craniectomy in the treatment of severe refractory posttraumatic cerebral edema. *Neurosurgery* 41:84–94, 1997
  72. Sahuquillo J, Arikan F: Decompressive craniectomy for the treatment of refractory high intracranial pressure in traumatic brain injury. *Cochrane Database Syst Rev* 25:CD003983, 2006
  73. Teasdale G, Parker L, Murray G, Knill-Jones R, Jennett B: Predicting the outcome of individual patients in the first week after severe head injury. *Acta Neurochir Suppl (Wien)* 28:161–164, 1979

